

An Assessment of The Knowledge, Risk Perception and Attitudes of Healthcare Workers in A Tertiary Health Facility in Southwest Nigeria to The Covid 19 Pandemic

Adeniyi MA¹, Olajide TG^{2,*}, Aderukuola B³, Popoola GO⁴, Olasehinde KO¹, Adegbiyi WA⁵

¹Department of Community Medicine, Federal Teaching Hospital, Ido-Ekiti, Ekiti state

²Department of ENT, Federal Teaching Hospital, Afe-Babalola University College of Medicine and Health Sciences, Ekiti, Nigeria

³Department of Obstetrics and Gynaecology, Federal Teaching Hospital, Ido-Ekiti, Ekiti state

⁴Department of Mental Health, Federal Teaching Hospital, Ido-Ekiti, Ekiti state

⁵Department of ENT, Ekiti State University Teaching Hospital, Ado Ekiti, Nigeria

Corresponding author:

Olajide Toye Gabriel, Department of Ear, Nose and Throat, Afe Babalola University, Ado Ekiti and Federal Teaching Hospital, Ido Ekiti, Nigeria

Keywords:

Knowledge, risk perception, attitude, health care workers

Received: Aug 26, 2022

Accepted: Oct 03, 2022

Published: Oct 18, 2022

Editor:

Ian James Martin, Edith Cowan University, USA

DOI: 10.14302/issn.2641-4538.jphi-22-4294

Abstract

Aim: Covid 19 is a pandemic that has ravaged the world resulting in thousands of deaths. This study aims to assess the knowledge, risk perception, and attitude of health care workers (HCWs) to the pandemic.

Methodology: This was a hospital based descriptive

cross sectional study conducted among health care workers who were staff of the hospital. Data collected includes socio demographic characteristics, data on knowledge, risk perception and attitude.

Results: A total of 288 HCWs participated in the study. Participants had good knowledge at 95.6%, major source of knowledge includes internet/social media 68.8%, radio 34.6% and television 28.3%. About 92.6% of the respondents also had positive attitude while 89.75 of the respondents had a positive risk perception.

Conclusion: Overall, there was a high level of knowledge, high positive attitude and a high risk perception in the study. Consequently, to ensure that this high level is maintained, there is need for continuous health education and promotion.

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARSCoV-2), is the pathogen that causes coronavirus disease2019 (COVID-19), and is one of the most contagious viruses in human history^{1,2}. It is a highly communicable viral respiratory infectious

disease with potential life-threatening complications.^{3,4,5} It is a single-stranded RNA virus that belongs to the family of coronaviridae and it is commonly transmitted through contact with infected respiratory droplets.^{3,6,7} The incubation period of COVID-19 is 1 – 14 days while its common symptoms include fever, cough, dyspnoea and headache.^{1,3,4} In addition, severe COVID-19 disease can lead to serious complications such as encephalitis, acute myocardial infection, renal failure, and multi-organ failure.^{1,2,4,7}

There is no specific treatment for this disease, so health care providers treat the clinical symptoms (e.g. fever, difficulty in breathing) which the patients present with. Supportive care (e.g. fluid management, oxygen therapy, etc.) can be highly effective for patients with symptoms.⁸

The first reported case of Covid-19 in Nigeria was in Lagos and this was reported by the Federal Ministry of Health on the 27th of February, 2020.¹⁰ As at 8th June, 2020 over 800 healthcare workers in Nigeria has been infected with the disease.¹¹ As at 16th of September, 2020 there are 56604 confirmed cases with 1091 deaths.⁹ Since the outbreak the Nigerian government, like other global community adopted measures to contain the spread of the disease. Some of these measures include social distancing, ban on public gathering including religious gatherings, continuous personal hygiene such as hand washing and use of hand sanitizers; use of face masks, limiting number of passengers in public vehicles, locking down public places etc.¹⁸ These measures have not been effective due partly to lack of government enforcement, poor knowledge and attitude of the citizen.

Statement of Problem

Globally, around 14% of COVID-19 cases reported to WHO are among health workers, and in some countries it's as much as 35%. The WHO estimates that over 41,000 health workers in Africa have been infected with COVID-19 as at 17/09/2020.¹⁹

Benefit

It is hoped that this study will help to improve the

knowledge of HCWs, make HCWs more aware and have a better understanding of the risks inherent in the Covid 19 pandemic and thereby improve their attitude and with an improved attitude reduce the risk of infection with Covid 19.

It is also hoped that this research will enable policy maker make informed and evidence based policies to better the lot of HCWs in the hospitals and the communities at large.

Furthermore, a better knowledge, a better understanding of the risks and a good attitude will make adherence to the preventive protocols put in place by the government and respective hospital managements easy to adhere to and follow.

Objectives

This study aims to assess the knowledge risk perception and attitude of healthcare workers in a tertiary health facility to the Covid 19 pandemic

Methodology

This is a descriptive cross sectional study carried out in a tertiary health centre located in Ido – Osi Local Government Area of Ekiti State. The study targets HCWs who are staff of the facility.

Ekiti State is located in the South-western part of Nigeria. Ido Ekiti town is the capital of Ido – Osi LGA which is one of the 16 LGAs in Ekiti State. It is also a semi – urban LGA.¹² Ido-Osi LGA lies within latitude 7046.715'N and 7055.822'N and longitude 508.410'E and 5014.416'E.¹³ The study was carried out in the Federal Teaching Hospital Ido-Ekiti which is one of the three tertiary health facilities located within the state. The other two tertiary health centres are located in the state capital Ado – Ekiti.¹²

Ido – Osi local Government is basically agrarian with people cultivating root crops and grains such as yams, cassava, maize, rice etc. There are a few cottage industries in the local Government devoted primarily to processing of Agricultural produce. Other industrial undertaking is in the area of printing, bakeries, weaving, carpentry etc.¹²

Sample Size Determination

$$n = Z^2 \times P (1 - P) / E^2$$

Where Z standard normal deviate at 95% = 1.96

P proportion of HCW with good knowledge in a similar Study = 78.6¹⁵

E Level of error = 5%

$$n = \frac{(1.96)^2 \times 78.6 (100 - 78.6)}{5^2}$$

$$= \frac{3.842 \times 78.6 (21.4)}{25}$$

$$= 258.5$$

$$= 259$$

$$= 259$$

$$= 259$$

For non-response rate compensation

$$ns = n / 0.9$$

$$ns = 259 / 0.9$$

$$= 288$$

Therefore, 288 HCWs participated in the study

Sampling Technique

All consenting HCWs in the tertiary health facility were recruited for the study until the sample size was met. Self-Administered Questionnaires were administered on them within the hospital premises.

Sampling Instrument / Tool

Data was collected using pre tested semi-structured self-administered questionnaires which was administered by the researchers and their trained assistants.

Pre testing of the questionnaire was done at the Ekiti State University Teaching Hospital (EKSUTH) located at Ado Ekiti. Which is also a tertiary health care facility.

Study Variables

Independent Variable

Socio demographic characteristics

Dependent Variables

knowledge, risk perception and attitude.

Data Analysis

Filled questionnaires were sorted and data was analysed using Statistical Package for Social Sciences (SPSS) version 22. Data were then presented in the form of frequency tables, cross tabulation using chi-square. Bivariate analysis was done as applicable. Level of significance was set with P-value less than 0.05.

Inclusion And Exclusion Criteria

Inclusion Criteria

All consenting health care workers who are staff of the hospital.

Exclusion Criteria

Patients, visitors and all non-staff were excluded.

Ethical Consideration: Ethical approval for this research was gotten from the Human Research and Ethics Committee of the Federal Teaching Hospital, Ido-Ekiti. Verbal Informed consent was gotten from all the respondents and only those who consented were recruited for the study.

The study was carried out between August 2020 and April 2021.

Results

Table 1 shows the Socio demographic characteristics of the study participants. The total number of HCWs surveyed was 272. Among them, 118 (43.4%) were between the age of 30-39 years, the mean \pm SD is 37.78 ± 8.17 with a range of 20 – 58.

Most of the respondents surveyed were female 164 (60.3%) while 118 (69.1%) had tertiary education qualification. Similarly, most of the HCW surveyed were nurses 88 (32.4%)

Table 2 and Figure 2 most of the respondents surveyed 260(95.6%) had good knowledge of Covid 19

Table 3 and Figure 3 showed that most of the respondents surveyed had positive attitude 252 (92.6%) while Table 4 and Figure 4 showed that risk perception among most of the respondents were positive 244 (89.7%).

Table 1. Socio-demographic distribution of the study participants

Variable	Frequency	Percent
Age (years)		
20 – 29	41	15.1
30 – 39	118	43.4
40 – 49	81	29.8
50 – 59	32	11.8
Mean ± SD	37.78 ± 8.17	
Range	20 – 58	
Sex		
Male	108	39.7
Female	164	60.3
Educational qualification		
Secondary	21	7.7
Tertiary	188	69.1
Post Tertiary	63	23.2
Marital status		
Single	44	16.2
Married	222	81.6
Divorced	3	1.1
Widow/ widower	3	1.1
Religion		
Christianity	262	96.3
Islam	7	2.6
Traditional	3	1.1
Tribe		
Yoruba	248	91.2
Ibo	10	3.7
Hausa	5	1.8
Others	9	3.3
Cadre		
Doctors	73	26.8
Nurses	88	32.4
Pharm	6	2.2
Lab	14	5.1
CHEW	36	13.2
Health attendants	41	15.1
Others	14	5.1
Years of practice		
< 5 Years	58	21.3
5 - 10 Years	81	29.8
> 10 Years	133	48.9

Table 2. Knowledge of HCWs on Covid 19

Knowledge of COVID 19	n (%)	n (%)
Have you ever heard of the COVID 19 is 1 to 14	272(100)	0(0.0)
Do you know that the COVID 19 pandemic is caused by SARS cov 2	186(68.4)	86(31.6)
Do you know that the COVID 19 pandemic was first reported in Wuhan China	239(87.9)	33(12.1)
The incubation period of COVID 19 is 1 to 14 days	253(93..0)	19(7.0)
The virus is spread mainly from person to person	254(93.4)	18(6.6)
The virus is also airborne	234(86.0)	38(14.0)
The following are some of the symptoms of COVID 19- cough, fever and difficulty in breathing	260(95.6)	12(4.4)
There is no known drug for COVID 19	223(82.0)	49(18.0)
Use of hand sanitizer and face mask can prevent the infection	262(96.3)	10(3.7)
Regular hand washing and physical social distancing can help prevent the spread of the disease	266(97.8)	6(2.2)
Presence of underlying illness such as hypertension and diabetes can make the infection more severe	248(91.2)	24(8.8)

Table 3. Attitude of HCWs to Covid 19

	SA	A	N	SD	D
Attitude	n (%)	n (%)	n (%)	n (%)	n (%)
Do you agree that COVID 19 will eventually be successfully controlled	123(45.2)	134(49.3)	10(3.7)	2(0.7)	3(1.1)
In your opinion will good personal hygiene reduce the risk of COVID 19 infection	106(39.0)	144(52.9)	12(4.4)	7(2.6)	3(1.1)
Do you agree that guidelines from the hospital COVID 19 taskforce will help in controlling the disease in the hospital	113(41.5)	123(45.2)	22(8.1)	4(1.5)	10(3.7)
Will you work or live with people who have been infected with COVID 19	29(10.7)	69(25.4)	55(20.2)	51(18.8)	68(25.0)
People infected with COVID 19 are careless with their health	22(8.1)	36(13.2)	29(10.7)	87(32.0)	98(36.0)
COVID 19 infected individuals should be isolated even after treatment	20(7.4)	41(15.1)	32(11.8)	77(28.3)	102(37.5)
COVID 19 can be transmitted through sexual intercourse	34(12.5)	56(20.6)	30(11.0)	67(24.6)	85(31.3)
People infected with COVID 19 should not be allowed to come to work	113(41.5)	99(36.4)	29(10.7)	12(4.4)	19(7.0)
Tuberculosis/Leprosy is better than COVID 19	31(11.4)	67(24.6)	54(19.9)	56(20.6)	64(23.5)
COVID 19 is a punishment from God	28(10.3)	26(9.6)	35(12.9)	120(44.1)	63(23.2)

Table 4. Risk perceptions of HCWs to Covid 19

Risk perceptions	n (%)	n (%)
COVID 19 is a severe debilitating infection	237(87.1)	35(12.9)
COVID 19 is highly infectious	257(94.5)	15(5.5)
COVID 19 can cause respiratory failure	251(92.3)	21(7.7)
COVID 19 can cause severe diarrhea	161(59.2)	111(40.8)
COVID 19 can cause severe dehydration	199(73.2)	73(26.8)
COVID 19 can cause lung injury for those who recover	174(64.0)	98(36.0)
It is safe to see a COVID 19 patient with PPE	235(86.4)	37(13.6)
It is safe to operate on a COVID 19 patient without PPE	58(21.3)	214(78.7)
It is safe to live in the same house with a COVID 19 patient without maintaining physical distance	27(9.9)	245(90.1)
It is safe to stay in the same office with an asymptomatic COVID 19 patient with the AC on	30(11.0)	242(89.0)
There is no risk of infection if you see a COVID 19 patient in the same clinic with other patients	55(20.2)	217(79.8)
Disinfecting the clinics and waiting area with hydrogen peroxide reduces the chances of infection to zero	132(48.5)	140(51.5)
Wearing PPE's (face mask, googles, gloves etc) will reduce the chances of infection among HCWs	251(92.3)	21(7.7)
Will you attend to a confirmed COVID 19 patient if provided with PPEs	231(84.9)	41(15.1)
Will you enter a ward where COVID 19 patients are kept if need be	168(61.8)	104(38.2)




Figure 1. Knowledge of COVID 19 among the healthcare workers




Figure 2. Source of knowledge about COVID 19




Figure 3. Attitude of health care workers to COVID 19




Figure 4. Overall risk perception of health workers to COVID 19

Table 4 and Figure 4 shows the risk perception of the HCWs to Covid 19. A greater percentage of the HCWs 244(89.3%) had a positive risk perception while only 28 (10.3%) had a negative perception.

Table 5 showed the association between socio demographic characteristics and attitude with marital status found to be significantly associated with attitude at a p-value of 0.020 with divorce and single respondents having the highest value of 100% and 97.7% respectively. Similarly, the cadre of the respondents was also found to be significantly associated with attitude at a p value of 0.003 with doctors and laboratory scientist having a value of 100% each.

Table 6 showed the association between socio demographic characteristics and risk perception. This study found sex to be of significant association with risk

perception at a p value of 0.013 with males having a higher positive risk perception 103 (95.4%) than females 141 (86%). Also, years of practice was found to be significantly associated with risk perception at a p value of 0.009 with those who have been in practice for more than 10 years having the highest value of positive risk perception at 122 (91.7%).

Table 7 shows the association between socio demographic characteristics and knowledge. This study found that 17(81%) , 180(95.7%) and 63(100%) of the respondents with secondary, tertiary and post tertiary education respectively had good knowledge of Covid 19 at a significant p value of 0.005. Similarly, there was also a significant association between years of practice and knowledge at a p value of 0.038 where 52 (91.4%), 76 (93%) and 131(98.5%) of the respondents with <5 years,

Table 5. Association between socio-demographic characteristics and attitude

	Positive	Negative	Total	χ^2	<i>p</i> value
Variables	n (%)	n (%)	N (%)		
Age					
20 – 29	39 (95.1)	2 (4.9)	41	4.300 ^F	0.211
30 – 39	109 (92.4)	9 (7.6)	118		
40 – 49	72 (88.9)	9 (11.1)	81		
50 – 59	32 (100.0)	0 (0.0)	32		
Sex					
Male	100 (92.6)	8 (7.4)	108	0.001	0.978
Female	152 (92.7)	12 (7.3)	164		
Educational qualification					
Secondary	17 (81.0)	4 (19.0)	21	5.214 ^F	0.059
Tertiary	174 (92.6)	14 (7.4)	188		
Post tertiary	61 (96.8)	2 (3.2)	63		
Marital status					
Single	43 (97.7)	1 (2.3)	44	9.707 ^F	0.020*
Married	205 (92.3)	17 (7.7)	222		
Divorced	3 (100.0)	0 (0.0)	3		
Widow/ widower	1 (33.3)	2 (66.7)	3		
Religion					
Christianity	242 (92.4)	20 (7.6)	262	0.244 ^F	1.000
Islam	7 (100.0)	0 (0.0)	7		
Traditional	3 (100.0)	0 (0.0)	3		
Tribe					
Yoruba	229 (92.3)	19 (7.7)	248	0.392 ^F	1.000
Others	23 (95.8)	1 (4.2)	24		
Cadre					
Doctor	73 (100.0)	0 (0.0)	73	17.488 ^F	0.003*
Nurses	82 (93.2)	6 (6.8)	88		
Pharm	5 (83.3)	1 (16.7)	6		
Lab	14 (100.0)	0 (0.0)	14		
CHEW	32 (88.9)	4 (11.1)	36		
Health attendants	35 (85.4)	6 (14.6)	41		
Others	11 (78.6)	3 (21.4)	14		
Years of practice					
< 5 years	52 (89.7)	6 (10.3)	58	1.067 ^F	0.623
5 – 10 years	76 (93.8)	5 (6.2)	81		
>10 years	124 (93.2)	9 (6.8)	133		

χ^2 : Chi square test; F: Fisher's exact test; *: *p* value <0.05

Table 6. Association between socio-demographic characteristics and risk perception

	Good	Poor	Total	χ^2	<i>p</i> value
Variables	n (%)	n (%)	N (%)		
Age					
20 – 29	37 (90.2)	4 (9.8)	41	1.294 ^F	0.754
30 – 39	103 (87.3)	15 (12.7)	118		
40 – 49	74 (91.4)	7 (8.6)	81		
50 – 59	30 (93.8)	2 (6.3)	32		
Sex					
Male	103 (95.4)	5 (4.6)	108	6.224	0.013*
Female	141 (86.0)	23 (14.0)	164		
Educational qualification					
Secondary	19 (90.5)	2 (9.5)	21	4.715	0.095
Tertiary	164 (87.2)	24 (12.8)	188		
Post tertiary	61 (96.8)	2 (3.2)	63		
Marital status					
Single	42 (95.5)	2 (4.5)	44	2.049 ^F	0.580
Married	196 (88.3)	26 (11.7)	222		
Divorced	3 (100.0)	0 (0.0)	3		
Widow/ widower	3 (100.0)	0 (0.0)	3		
Religion					
Christianity	234 (89.3)	28 (10.7)	262	0.293 ^F	1.000
Islam	7 (100.0)	0 (0.0)	7		
Traditional	3 (100.0)	0 (0.0)	3		
Tribe					
Yoruba	221 (89.1)	27 (10.9)	248	1.070 ^F	0.486
Others	23 (95.8)	1 (4.2)	24		
Cadre					
Doctor	69 (94.5)	4 (55.5)	73	7.867 ^F	0.192
Nurses	76 (86.4)	12 (13.6)	88		
Pharm	4 (66.7)	2 (33.3)	6		
Lab	14 (100.0)	0 (0.0)	14		
CHEW	31 (86.1)	5 (13.9)	36		
Health attendants	37 (90.2)	4 (9.8)	41		
Others	13 (92.9)	1 (7.1)	14		
Years of practice					
< 5 years	56 (96.6)	2 (3.4)	58	9.466 ^F	0.009*
5 – 10 years	66 (81.5)	15 (18.5)	81		
>10 years	122 (91.7)	11 (8.3)	133		

χ^2 : Chi square test; F: Fisher's exact test; *: *p* value <0.05

Table 7. Association between socio-demographic characteristics and knowledge

	Good	Poor	Total	χ^2	p value
Variables	n (%)	n (%)	N (%)		
Age					
20 – 29	39 (95.1)	2 (4.9)	41	0.428 ^F	0.971
30 – 39	112 (94.9)	6 (5.1)	118		
40 – 49	78 (96.3)	3 (3.7)	81		
50 – 59	31 (96.9)	1 (3.1)	32		
Sex					
Male	104 (96.3)	4 (3.7)	108	0.213 ^F	0.768
Female	156 (95.1)	8 (4.9)	164		
Educational qualification					
Secondary	17 (81.0)	4 (19.0)	21	10.245 ^F	0.005
Tertiary	180 (95.7)	8 (4.3)	188		
Post tertiary	63 (100.0)	0 (0.0)	63		
Marital status					
Single	42 (95.5)	2 (4.5)	44	1.141 ^F	1.000
Married	212 (95.5)	10 (4.5)	222		
Divorced	3 (100.0)	0 (0.0)	3		
Widow/ widower	3 (100.0)	0 (0.0)	3		
Religion					
Christianity	251 (95.8)	11 (4.2)	262	2.757 ^F	0.368
Islam	6 (85.7)	1 (14.3)	7		
Traditional	3 (100.0)	0 (0.0)	3		
Tribe					
Yoruba	237 (95.6)	11 (4.4)	248	0.004 ^F	1.000
Others	23 (95.8)	1 (4.2)	24		
Cadre					
Doctor	71 (97.3)	2 (2.7)	73	7.300 ^F	0.200
Nurses	86 (97.7)	2 (2.3)	88		
Pharm	6 (100.0)	0 (0.0)	6		
Lab	12 (85.7)	2 (14.3)	14		
CHEW	34 (94.4)	2 (5.6)	36		
Health attendants	37 (90.2)	4 (9.8)	41		
Others	14 (100.0)	0 (0.0)	14		
Years of practice					
< 5 years	53 (91.4)	5 (8.6)	58	6.036 ^F	0.038
5 – 10 years	76 (93.8)	5 (6.2)	81		
>10 years	131 (98.5)	2 (1.5)	133		

χ^2 : Chi square test; F: Fisher's exact test; *: p value <0.05

Table 8. Association between knowledge, attitude and risk perception

	Good	Poor	Total	χ^2	<i>p</i> value
Variables	n (%)	n (%)	N (%)		
Knowledge					
Good	236 (90.8)	24 (9.2)	260	7.216 ^F	0.025*
Poor	8 (66.7)	4 (33.3)	12		
Attitude					
Positive	225 (89.3)	27 (10.7)	252	0.655 ^F	0.704
Negative	19 (95.0)	1 (5.0)	20		

χ^2 : Chi square test; F: Fisher's exact test; *: *p* value <0.05

5-10 years and >10 years, years of experience post graduation had good knowledge showing a progressive increase in the percentage with increasing years of experience.

Table 8 shows the association between knowledge, attitude and risk perception. This study found that there was a significant association between knowledge and risk perception at a *p* value of 0.025.

Discussion

This study found that all the respondents have heard of COVID-19. This is not surprising considering that the study was carried out among HCW and also Covid 19 is a pandemic currently ravaging the world. Similar findings were also recorded in studies done in Nigeria, Uganda, China and other parts of the world^{15, 21-24} Generally, majority 260 (95.6%) of the respondents in this study had good knowledge of Covid 19, this is also similar to findings in studies done in Nigeria^{15,21} Greece,²⁵ China²³ and other parts of the world^{15,22, 26,27}

This study also found out that the most common source of information amongst the respondents on the disease was the internet/social media 187 (68.8%) followed by the Radio 94 (34.6%), this is in contrast to a study done in Greece²⁴ and part of Nigeria²⁴ where the Radio was the most common source of information followed by the internet/social media. However, the

findings were similar to those of studies done Nigeria and China^{20,23} consequently, considering the importance of information dissemination to the control and eradication of Covid 19 stakeholders should use the internet/social media to disseminate information about Covid 19 more as this is a major source of information for HCWs. The traditional method of dissemination of information such as the radio and television should nonetheless not be discarded as this study also found that they are still a useful source of information dissemination.

Attitudes of people as a mediator between their knowledge and practices have an important role for better controlling of epidemics and pandemics of infectious diseases; they facilitate the process of changing people's behaviour. Also, risk perceptions can influence health-related behaviours and change risky behaviours thus giving individuals a better outcome in the event of a health challenge.²⁸ This study found that overall; the attitude of the majority of the HCWs was positive as evidenced by a total number of 252 (92.6%) HCWs with a positive attitude. This positive attitude may also be connected to the good knowledge exhibited by the respondents who are HCWs in this study. Similarly, this good knowledge was also associated with a high level of positive risk perception 244 (89/7%), similar findings were reported in studies done Nigeria¹⁵ Greece²⁵ China²⁹

however, findings in this study contrasted sharply with findings in a study done in Iran where only about 50% of the respondents had a positive attitude towards Covid 19³⁰. Study done by Srichan et al³¹ revealed that knowledge and attitudes toward disease prevention and control are poor in northern Tanzanian, while another study in Peru showed that over 75% of the respondents had a poor attitude to covid 19³². Ilesanmi et al in their study showed that there no significant difference existed in the risk perception, and KAP towards COVID-19 prevention among Doctors and Nurses³³. Most of the respondents in this study 236 (86.7%) agreed to follow the guidelines of the hospital Covid 19 taskforce this is in sharp contrast to findings in a study done in Thailand where only 15% of the respondents agreed to follow the Covid 19 guidelines

There is yet no cure for Covid 19 despite the extensive research on going on the diseases. The fact still remains that Covid 19 is a serious life threatening disease with a high mortality rate particularly in severe cases.^{1,4,5} This study found that there was a high risk perception to covid 19, this may again be due to the fact that the sampling population were made up of HCWs. 244 (89.7%) of the respondents had a positive risk perception and this was similar to findings in another study done in Nigeria²⁰

Limitations and Recommendations

This study is a hospital based study where the knowledge, attitude and risk perception is expected to be high due to the fact that the study population is made up of HCWs and as such can be used to use to assess the level of knowledge of health personnel which is expected to be higher than that of the community. This may also be a limitation, a community based study will help take care of this limitation, and similarly a comparative study between urban and rural communities will be more revealing. We recommend that similar study be conducted among different classes of people such as teachers, bankers, artisans, clergy etc

Conclusion

Overall, there was a high level of knowledge, high

positive attitude and a high risk perception in the study. Consequently, to ensure that this high level is maintained, there is need for continuous health education and promotion.

Conflict of interest

None was declared

Funding

This study was self funded

Ethical Approval

Ethical approval was sought and gotten from the Human Research and Ethical committee of the Federal Teaching hospital, Ido- Ekiti. Ekiti state

References

1. World Health Organization, *Coronavirus Disease 2019 (COVID-19): Situation Report, World Health Organization, Geneva, Switzerland, 2020*
2. J. F.-W. Chan, K.-H. Kok, Z. Zhu et al., "Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan," *Emerging Microbes & Infections*. 2020;9(1):221-236
3. Yi Y, Lagniton PNP, Ye S, Li E, Xu RH. COVID-19: what has been learned and to be learned about the novel coronavirus disease. *Int J Biol Sci*.2020;16(10): 1753-1766. doi: 10.7150/ijbs.45134.
4. Carod-Artal FJ. Neurological complications of coronavirus and COVID-19. *Rev Neurol*.2020;70 (9):311-322. doi:10.33588/rn.7009.2020179
5. WorldHealthOrganization.Coronavirus.https://www.who.int/healthtopics/coronavirus#tab=tab_1 Accessed May 14, 2020.
6. Lai CC, Shih TP, Ko WC, Tang HJ, Hsueh PR. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The epidemic and the challenges. *Int JAntimicrob Agents*. 2020;55(3):105924.
7. Guo YR, Cao QD, Hong ZS, Tan YY, Chen SD, JinHJ, et al. The origin, transmission and clinicaltherapies on

- coronavirus disease 2019 (COVID-19) outbreak - an update on the status. *Mil Med Res.* 2020;7(1):11
8. Labban L, Thallaj N, Labban A (2020) Assessing the Level of Awareness and Knowledge of COVID 19 Pandemic among Syrians. *ArchMed* Vol.12 Iss.2: 8
 9. covid19: ncdc.gov.ng (Last accessed on 17/09/2020)
 10. First case of Corona virus disease (Covid 19) confirmed in Nigeria. Available at ncdc.gov.ng/news/227
 11. Nigeria: 800 health workers infected with Covid 19. Available at www.africanews.com/2020/09/15/nigeria-800-health-workers-infected-with-covid-19/ (last Accessed on 15/09/2020)
 12. Ekiti State Website - <https://ekitistate.gov.ng> (last accessed on 17/09/2020).
 13. Ido-Osi map - Nigeria - mapcarta available @ [mapcarta.com](https://www.mapcarta.com) (last accessed on 16/09/2020)
 14. Jekel JF, Katz DL, Elmore JG, Wild D. Sample size, randomization and probability theory. *Epidemiology, Biostatistics and preventive medicine.* Philadelphia Saunders; 2001(2):194-199.
 15. Olubunmi A, Aishat BU, Posi A, Abass G. Cross-sectional study on Oyo State Health Care workers Knowledge, Attitude and Practice regarding Corona Virus Disease 2019(COVID-19). Available at Research Square [[+https://doi.org/10.21203/rs.3.rs-23501/v1](https://doi.org/10.21203/rs.3.rs-23501/v1)]
 16. Covid 19 Treatment guidelines Panel. Coronavirus disease 2019 (COVID-19) treatment Guidelines. National Institute of Health. Available at <https://www.covid19treatmentguidelines.nih.gov/>. Accessed on 16/09/20
 17. National Interim Guidelines for Clinical Management of Covid 19. Nigeria Centre for Disease Control. Available at ncdc.gov.ng/covid19, ncdc.gov.ng. Accessed on 14/09/2020
 18. Nigeria Centre for Disease Control. (2020). Implementation guidelines for containment of Covid-19.
 19. WHO DG's opening remarks at the member state briefing on COVID-19. Available at <https://www.who.int/dg/speeches/details/who-director-general-s-opening-remarks-at-the-member-state-briefing-on-covid-19-17-september-2020>
 20. Kanmodi KK, Nwafor NJ, Amoo BA, Akinbami AA, Hundeji AA. Knowledge, risk perception, and attitudes of Nigerian dentists toward Covid-19: Pilot study. *Yen Med J.* 2020;2(4):30 – 37.
 21. Zhong BL, Luo W, Li HM. Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID- 19 outbreak: a quick online cross-sectional survey. *Int J Biol Sci.* 2020;16(10):1745-1752
 22. Olum R, Chekwech G, Wekha G, Nassozi DR, Bongomin F. Coronavirus Disease-2019: Knowledge, Attitude, and Practices of Health Care Workers at Makerere University Teaching Hospitals, Uganda. *Front Public Health.* 2020;8:181.
 23. Bhagavathula AS, Aldhaleei WA, Rahmani J, Mahabadi MA, Bandari DK. Knowledge and Perceptions of COVID -19 Among Health Care Workers: Cross-Sectional Study. *JMIR Public Health Surveill.* 2020;6(2):e19160
 24. Fadeyi A, Adeoti A.O, Adeboye MAN, Awosanya J.A, Oluwadiya I.O, Oluwadiya K.S. KNOWLEDGE, ATTITUDE AND RISK PERCEPTION OF COVID-19 AMONG NIGERIANS. *Fadeyi et al., Afr., J. Infect. Dis.* (2022) 16(2): 63-70
 25. Dimitrios P , Foteini M, Dimitrios GR, Ioanna VP , Evangelos C.F , Zoe D et al. Assessment of Knowledge, Attitudes, and Practices towards New Coronavirus (SARS-CoV-2) of Health Care Professionals in Greece before the Outbreak Period. *Int. J. Environ. Res. Public Health* 2020, 17, 4925; doi:10.3390/ijerph17144925
 26. Nwagbara UI, Osual EC, Chireshe R, Bolarinwa OA, Saeed BQ, Khuzwayo N, et al. (2021) Knowledge, attitude, perception, and preventative practices towards COVID-19 in sub-Saharan Africa: A scoping review. *PLoS ONE* 16(4): e0249853

27. Iheanacho T, Stefanovics E, Okoro UG, et al. Assessing knowledge, attitude, practice and training related to COVID-19: a cross-sectional survey of frontline healthcare workers in Nigeria. *BMJ Open* 2021;11:e050138
28. Behnam H, Kamran B. L, Ali K, Fatemeh S, Mohammadhassan Z, Mohammad Reza R H et al. Knowledge, attitudes, risk perceptions, and practices of adults toward COVID-19: a population and field-based study from Iran. *International Journal of Public Health* · June 2020 DOI: 10.1007/s00038-020-01406-2
29. Chen Y, Jin YL, Zhu LJ, Fang ZM, Wu N, Du MX et al. The network investigation on knowledge, attitude and practice about Novel coronavirus pneumonia of the residents in Anhui Province *Zhonghua yufang yi xue za zhi. Chinese journal of preventive medicine.* 2020; 17; 54(0): E004. <https://doi.org/10.3760/cma.j.issn.0253-9624.2020.0004>
30. Behnam H, Kamran B L, Ali K, Fatemeh S, Mohammadhassan Z, Mohammad Reza RH et al Knowledge, attitudes, risk perceptions, and practices of adults toward COVID-19: a population and field-based study from Iran. *International Journal of Public Health.* 2020. <https://doi.org/10.1007/s00038-020-01406-2>
31. Srichan P, Apidechkul T, Tamornpark R, Yeemard F, Khunthason S, et al. Knowledge, attitude and preparedness to respond to the 2019 novel coronavirus (COVID-19) among the bordered population of northern Thailand in the early period of the outbreak: a cross-sectional study. *WHO South East Asia journal of public health.* 2020; 992):118-125
32. Zegarra-Valdivia JA, Chino-Vilca BN, Ames-Guerrero R. Knowledge, perception and attitudes in Regard to COVID-19 Pandemic in Peruvian Population. *PsyArXiv [Internet].* 2020;(April). Available from: <https://www.researchgate.net/publication/340694053%0AKnowledge>.
33. Ilesanmi O.S., Kareem A.O, Afolabi A.A, Kareem A.J and Ukwanya V. RISK PERCEPTION, KNOWLEDGE, ATTITUDE AND PRACTICES TOWARDS COVID-19 AND LASSA FEVER PREVENTION AMONG DOCTORS AND NURSES IN A TREATMENT CENTRE IN NIGERIA. *Vol. 19 2021 COVID-19 Supplement S8 Ann Ibd. Pg. Med 2021. Vol.19 COVID-19 Supplement S8 - S14*